Local Data: Atlanta

In 2020, there were 38,140 people living with HIV in Atlanta. In 2020, 1,254 people were newly diagnosed with HIV.

HIV Prevalence

Number of people living with HIV, 2020


38,140

Rate of people living with HIV per 100,000 population, 2020

Aged 13-24

1,088

Percent of people living with HIV, by Sex, 2020




Aged 25-44

Rate of people living with HIV per 100,000 population, by Geography, 2020

Aged 60+

Aged 45-59



New HIV Diagnoses

Number of new HIV diagnoses, 2020


1,254

Rate of new HIV diagnoses per 100,000 population, 2020

Aged 13-24

36

Percent of people newly diagnosed with HIV, by Sex, 2020




Aged 25-44

Number of New HIV Diagnoses, 2016-2020


Aged 60+

Aged 45-59



Due to the COVID-19 pandemic, data from 2020 should be interpreted with caution

Rate of people newly diagnosed with HIV per 100,000 population, by Geography, 2020



Late HIV Diagnoses

A late HIV diagnosis is defined as having an AIDS diagnosis within three months of initial HIV diagnosis.

Number of new HIV diagnoses that were diagnosed late, 2020

258

Proportion of people with a late HIV diagnoses, by Sex, 2020

Male: 20.4%

Female: 21.2%

Proportion of people with a late HIV diagnoses, by Race/Ethnicity, 2020

Black: 19.6%

Hispanic/Latinx: 23.5%

White: 21.0%

Percent of new HIV diagnoses that were diagnosed late, 2020

20.6%

Proportion of people with a late HIV diagnoses, by Age, 2020

Aged 13-24: 8.3%

Aged 25-44: 20.8%

Aged 45-59: 35.1%

Aged 60+: 32.4%

Linked to HIV Care

Linkage to care is defined as having a visit with an HIV health care provider within one month of being diagnosed with HIV.

Number of people diagnosed with HIV and linked to HIV care, 2020

962

Proportion of people linked to HIV care, by Sex, 2020

Male: 77.2%

Female: 74.5%

Proportion of people linked to HIV care, by Race/Ethnicity, 2020

Black: **75.8**%

Hispanic/Latinx: 77.2%

White: 80.3%

Percent of people diagnosed with HIV and linked to HIV care, 2020

76.7%

Proportion of people linked to HIV care, by Age, 2020

Aged 13-24: 67.3%

Aged 25-44: 79.6%

Aged 45-59: 77.3%

Aged 60+: 83.8%

Receipt of HIV Care

Receipt of HIV care is defined as those living with diagnosed HIV who received medical care for HIV and had at least one CD4 count or HIV viral load test in that year.

Number of people living with HIV who received HIV care, 2020

26,137

Proportion of people who received HIV care, by Sex, 2020

Male: 74.4%

Female: 71.7%

Proportion of people who received HIV care, by Race/Ethnicity, 2020

Black: **73.2**%

Hispanic/Latinx: 70.9%

White: **76.9%**

Percent of people living with HIV who received HIV care, 2020

73.9%

Proportion of people who received HIV care, by Age, 2020

Aged 13-24: 80.4%

Aged 25-44: 75.1%

Aged 45-59: 73.6%

Aged 60+: 70.0%

Viral Suppression

Viral suppression is defined as those living with diagnosed HIV who had suppressed HIV viral load (<200 copies/mL).

Number of people living with HIV who were virally suppressed, 2020

22,307

Proportion of people who were virally suppressed, by Sex, 2020

Male: 59.0%

Female: 56.3%

Proportion of people who were virally suppressed, by Race/Ethnicity, 2020

Black: **56.2**%

Hispanic/Latinx: 58.5%

White: 66.0%

Percent of people living with HIV who were virally suppressed, 2020

58.5%

Proportion of people who were virally suppressed, by Age, 2020

Aged 13-24: 47.8%

Aged 25-44: 55.9%

Aged 45-59: **61.8%**

Aged 60+: 60.6%

HIV Mortality

Number of deaths of people living with HIV in Atlanta, 2020

426

Number of deaths of people living with HIV in Georgia, 2019

806

Demographics, 2010


Total Population

4,333,441

City Population by Race/Ethnicity


- American Indian / Alaska Native (0.2%)
- Asian (5.5%)
- Black (34.6%)
- Hispanic / Latinx (11.5%)

- White (46.1%)
- Multiple Race (1.8%)
- Other* (0.3%)




^{*}Includes other races/ethnicities or missing/suppressed data


HIV Prevalence Rate Ratios, by Race/Ethnicity, 2020




Late HIV Diagnoses, 2020

A late HIV diagnosis is defined as having an AIDS diagnosis within three months of initial HIV diagnosis.






Region



United States




Diagnosed with HIV Diagnosed with AIDS

People Living with HIV, by Transmission Category, 2020


Percent of People Living with HIV, by Transmission Category, 2020





- Injection Drug Use (2.3%)
- Heterosexual Contact (4.1%)
- Male-to-Male Sexual Contact (72.7%)
- Male-to-Male Sexual Contact & Injection Drug Use (3.9%)
- Other* (17.0%)

Female Transmission Categories




^{*}Includes risk factor not reported or identified, along with hemophilia, blood transfusion, perinatal exposure, or missing/suppressed data.

People Newly Diagnosed with HIV, by Transmission Category, 2020


Percent of People Newly Diagnosed with HIV, by Transmission Category, 2020

Male Transmission Categories



- Injection Drug Use (0.8%)
- Heterosexual Contact (7.1%)
- Male-to-Male Sexual Contact (76.9%)
- Male-to-Male Sexual Contact & Injection Drug Use (1.6%)
- Other* (13.6%)


Female Transmission Categories




^{*}Includes risk factor not reported or identified, along with hemophilia, blood transfusion, perinatal exposure, or missing/suppressed data.

Social Determinants of Health, 2020






Percent of Population with a High School Education, 2020




Income Inequality (Gini Coefficient), 2020




Percent of Population Living in Poverty, 2020



Median Household Income, 2020



Sexually Transmitted Diseases, 2019



Rates of Sexually Transmitted Diseases per 100,000 Population, 2019

CityStateRegionUnited States

